

University of Groningen

Zirex

Mu, Dongdong; Montalbán-López, Manuel; Masuda, Yoshimitsu; Kuipers, Oscar P.

Published in:
Applied and environmental microbiology

DOI:
[10.1128/AEM.00866-13](https://doi.org/10.1128/AEM.00866-13)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2013

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Mu, D., Montalbán-López, M., Masuda, Y., & Kuipers, O. P. (2013). Zirex: a Novel Zinc-Regulated Expression System for *Lactococcus lactis*. *Applied and environmental microbiology*, 79(14), 4503-4508. <https://doi.org/10.1128/AEM.00866-13>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Dongdong Mu,^a Manuel Montalbán-López,^a Yoshimitsu Masuda,^a Oscar P. Kuipers^{a,b}Groningen-Delft, The Netherlands^b

Here, we report a new zinc-inducible expression system for *Lactococcus lactis*, called Zirex, consisting of the pneumococcal repressor SczA and P_{czcD}. P_{czcD} tightly regulates the expression of green fluorescent protein in *L. lactis*. We show the applicability of Zirex together with the nisin-controlled expression system, enabling simultaneous but independent regulation of different genes.

Lactococcus lactis is a Gram-positive bacterium that has been intensively engineered for the production of heterologous proteins (1, 2). In addition, it is an organism generally recognized as safe (GRAS). To date, several promoters originally from *Lactococcus*, regulated by inducers or environmental factors, have been documented, including the *dnaJ* promoter, induced by heat shock (3); the PA170 promoter, which can be upregulated at a low pH during the transition to stationary phase (4); the *prtP* promoter, which is regulated by the peptide concentration in the medium (5); and the P_{Zn} *zitR* promoter, which responds to divalent cation starvation (6). The promoter of nisin, P_{nisA} , is the most widely used promoter for inducible protein expression in *L. lactis* (1, 7) and other Gram-positive bacteria (8). The expression from the P_{nisA} promoter is regulated by the two-component regulatory system NisRK, which is triggered by nisin. For the other promoters

mentioned above, there are still some drawbacks, such as relatively low induction levels or high background level at the uninduced stage, which may complicate efforts to tightly control the expression or coexpression of one or two different proteins in the same cell (9). The aims of the present work were to develop a novel tightly controlled promoter for *L. lactis* and to investigate if such

Published ahead of print 10 May 2013

Published ahead of print 10 May 2013

Address correspondence to Oscar P. Kuipers, o.p.kuipers@rug.nl.

Copyright © 2013, American Society for Microbiology. All Rights Reserved.

doi:10.1128/AEM.00866-13




FIG 1 (A) Nucleic acid sequence of *P_{czcD}*. Motif1 (in boxes) and Motif2 (red letters) are the binding sites for the SczA regulator. The start codon of *czcD* is indicated in italics. The ribosome binding site (RBS) of *P_{czcD}* is shown in boldface. -35 and -10 sequences are underlined (10). (B) Expression systems used in this study. Hairpins represent terminators.

TABLE 1 Primers used in this study

Primer	Sequence ^a	Restriction site(s)
czcD-f	CGGGGTACCGGATCCTGCAGGCAGATATAGTTGATAATCAAGG	KpnI, SbfI
czcD-r	CAGCTCTTCTCCTTTTCCCATGGTTCTCATTCCCTTTGTTATAATAG	NcoI
gfp-f	CTATTATAACAAAGGAATGAGAACCATGGGAAAAGGAGAAGAGCTG	NcoI
gfp-r	GGAAGATCTATTAATCGAAATACGGGCAGAC	BglII
gfp-r2	CCCAAGCTTCGAAATACGGGCAGAC	HindIII
mCherry-f	CGGGGTACCTCCTGGTTGCAAATTTTG	KpnI
mCherry-r	CGTACTCACGTGCTGCAAGGCGATTAAGTTG	PmlI
sczA-czcD-f	ATCAAGATCTAGAAATAAGACAACCTGAAGCTTTAC	BglII
sczA-czcD-r	AGATCCATGGTTCTCATTCCCTTTGTTATAATAG	NcoI
pIL-f	ATCAAGATCTACAGCAAAGAATGGCGGAAACG	BglII
pIL-r	AATCGATAAGCTTGGCTGCAGGTC	

^a Restriction sites engineered in the primers are underlined.

an inducible promoter system could be coupled to the P_{nisA} promoter to create a dual-promoter-regulated production system for different proteins. First, we searched in the genome of *L. lactis* MG1363 (NCBI reference sequence [NC_009004.1](#)) for proteins putatively involved in cation transport that may be regulated by the presence of cations. A putative promoter, namely, P_{Zn3} , preceding the translation of a cationic ion efflux protein (NCBI reference sequence [YP_001032214.1](#)) in *L. lactis* was further investi-

gated (see below). Additionally, we explored the genome of other related Gram-positive bacteria for cation-regulated promoters. In the case of *Streptococcus pneumoniae*, a zinc-inducible promoter was previously described by Kloosterman et al. (10) and Eberhardt et al. (11). *sczA* and *czcD* are transcribed divergently (Fig. 1A). The promoter of *czcD* gene is regulated by SczA. SczA binding to the motif 2 sequence located downstream of the -10 sequence of P_{czcD} blocks transcription of *czcD* in the absence of zinc. After the




FIG 2 Growth (dotted lines) and GFP expression level (solid lines) after induction with 0.5 mM zinc at different time points. NZ9000(pCZG) induced with ZnSO_4 at a final concentration of 0.5 mM at 0 h (\square), 2 h (\triangle), and 4 h (\circ). Two control strains were run in parallel: NZ9000(pNZ8048) uninduced (\diamond) and induced with 0.5 mM ZnSO_4 at 0 h (\times). The values represent the means from three independent measurements.




FIG 3 (A) Growth (dotted lines) and GFP intensity (solid lines) after induction with different zinc concentrations. *L. lactis* NZ9000(pCZG) was induced at an OD₆₀₀ of 0.06 with different concentrations of ZnSO₄: 0 (◇), 0.1 mM (□), 0.7 mM (△), or 1.0 mM (○). A control, NZ9000(pNZ8048G) (×), induced with 5 ng/ml of nisin was also used to compare the production levels of P_{czcD} and P_{nisA}. The arrow indicates the time point for induction. These values represent the means from three independent measurements. (B) Dose-response curve of GFP expression of *L. lactis* NZ9000(pCZG). Fluorescent signal is shown as specific units per OD₆₀₀. The fluorescent signal changed less sensitively at the concentration of ZnSO₄ above 0.3 mM. The standard errors are less than 15% for each value.

addition of zinc, SczA will move to motif 1 unblocking the transcription (10).

Primers *czcD*-f and *czcD*-r (Table 1) were designed to amplify the regulator protein SczA and the P_{czcD} region from the *S. pneumoniae* D39 (12) genome, including the restriction sites KpnI and NcoI, respectively. The gene coding for green fluorescent protein

(GFP) with its own terminator was amplified from pJWV102_gfp (a kind gift from J. W. Veening) by PCR with primers *gfp*-f and *gfp*-r. A BglII site was added on the 5' end of primer *gfp*-r. *czcD*-r and *gfp*-f were designed to be reverse complementary by overlapping the 5' ends of each other, and an NcoI site was inserted in both primers. The fragment SczA-P_{czcD}-GFP was generated by

spliced overlap extension PCR with primers *czcD*-f and *gfp*-r using the mixture of *SczA*-*P_{czcD}* and GFP-specifying amplicons as the templates (13). After digestion with KpnI and BglII, *SczA*-*P_{czcD}*-GFP was cloned into pNZ8048 (7), digested with the same enzymes to create the plasmid pCZG (Fig. 1B). pZn3G was constructed based on pCZG, in which the *gfp* gene was controlled by *P_{Zn3}* (Fig. 1B). Unfortunately, the low production level obtained after induction with zinc and the leakage in the noninduced state made *P_{Zn3}* an unsuitable candidate for further characterization (data not shown). pNZ8048G was created by cloning *gfp* amplified with the primers *gfp*-f and *gfp*-r2 (Table 1) in the NcoI-HindIII sites of pNZ8048. In pNZ8048G, GFP expression is under the control of *P_{nisA}* (Fig. 1B).

The expression assays were carried out in *L. lactis* NZ9000 (9), which was transformed with pCZG (containing *SczA*-*P_{czcD}*-GFP) according to Holo and Nes (14). All of the expression assays were conducted at 30°C in a chemically defined medium for prolonged cultivation (CDMPC) without ZnSO₄ supplemented with 10 µg/ml chloramphenicol (B. Teusink, F. Santos, O. P. Kuipers, C. E. Price, J. Kok, and D. Molenaar, unpublished data). Each assay was repeated in triplicate in a 96-well microtiter plate and monitored with an Infinite 200 Pro microplate spectrophotometer (Tecan Group, Ltd., Mannedorf, Switzerland).

First, we investigated the optimal induction moment. For this purpose, ZnSO₄ was added after 0 h, 2 h, or 4 h of growth at a final concentration of 0.5 mM. The cell growth was monitored measuring the optical density at 600 nm (OD₆₀₀), and the signal of GFP was measured using an excitation wavelength of 485 nm and an emission wavelength of 535 nm (15). NZ9000(pNZ8048) was used as a negative control. We observed that the earlier we induced, the stronger the displayed signal was, with no induction observed in the stationary phase (Fig. 2). It should be noted that 0.5 mM ZnSO₄ showed comparatively low toxicity when it was introduced before inoculation (0 h) of the strains with (pCZG) or without (pNZ8048) GFP. The addition after 2 h of growth does not cause a visible reduction in growth. To assess if the slower growth when the cells were induced at 0 h was caused by the Zn salt used in the study, we also studied the growth tendencies and fluorescent signals of cells induced with the same amount of Zn²⁺ using ZnCl₂. The growth curves and the expression profiles were almost the same as those after induction with ZnSO₄ (data not shown). These results indicate that different Zn²⁺ sources do not affect the toxicity or potency of the induction.

In order to assess the optimal concentration of ZnSO₄ for the induction, NZ9000(pCZG) was induced at an OD₆₀₀ of 0.06 (the middle of the exponential phase) with a final concentration of 0, 0.1, 0.3, 0.5, 0.7, or 1.0 mM ZnSO₄. NZ9000(pNZ8048G) grown in CDMPC was induced with 5 ng/ml of nisin at an OD₆₀₀ of 0.06 as a reference. The growth rate was not affected by the addition of ZnSO₄, and the GFP signal produced by NZ9000(pNZ8048G) or NZ9000(pCZG) reached the highest level after 2 h or 2.5 h of induction, respectively (Fig. 3A). The highest intensity of GFP produced using the zinc-inducible system was almost 80% of that produced with the nisin-inducible system. Moreover, the GFP signal in the induced cells increased nearly proportionally with the ZnSO₄ concentration in the range between 0 and 0.3 mM (Fig. 3B). Furthermore, almost no fluorescent signal was detected under uninduced conditions, which demonstrated that this pneumococcal system is also effectively repressed in the absence of zinc in *L. lactis*.

TABLE 2 Simultaneous production of GFP and mCherry

ZnSO ₄ as inducer (0.7 mM) ^a	Nisin as inducer (5 ng/ml 1 h later) ^a	Fluorescent intensity (AU) ^b	
		GFP (2.5 h)	mCherry (2.5 h)
+	–	181.00 ± 8.01	0.67 ± 4.19
+	+	138.00 ± 2.49	544.33 ± 17.68
–	+	5.33 ± 3.00	612.00 ± 39.04

^a +, inducer present; –, inducer absent.

^b AU, arbitrary units.

pCZGM was constructed to observe the effect of the induction with nisin and zinc at the same time. In pCZGM, *P_{nisA}* controls the expression of mCherry, whereas *P_{czcD}* controls the expression of GFP. To construct this vector, a fragment encompassing from *P_{nisA}* to the terminator of mCherry was amplified from pHK35C (a generous gift from H. Karsens) using the primers mCherry-f and mCherry-r, containing at their 5' end a KpnI site and a PmlI site, respectively. After digestion with KpnI and PmlI, the fragment was inserted into pCZG cut with the same enzymes, resulting in pCZGM (Fig. 1B). The signal of mCherry was measured using an excitation wavelength of 590 nm and an emission wavelength of 620 nm, and GFP was measured as mentioned above. Cultures were induced with 0.7 mM ZnSO₄ at an OD₆₀₀ of 0.06 and with 5 ng/ml nisin 1 h later. Uninduced controls lacking either nisin or zinc were run in parallel. In Table 2, we can observe the expression level of GFP or mCherry achieved after 2.5 h of the induction with ZnSO₄ or nisin. These data show that simultaneous overexpressions of mCherry and GFP in this system cause around 23% and 11% reduction of the two fluorescent signals, respectively.

Based on the results described above, we created pCZ-Cm for general use as a chloramphenicol-resistant expression vector for *L. lactis*. In this vector, the multiple-cloning site (MCS) of pNZ8048 was fused behind *P_{czcD}*. For this purpose, the region *SczA*-*P_{czcD}* was amplified from pCZG with the primers *sczA*-*czcD*-f and *sczA*-*czcD*-r (Table 1). After digestion with BglII and NcoI, the fragment *SczA*-*P_{czcD}* was inserted into pNZ8048 digested with the same restriction enzymes, rendering pCZ-Cm (Fig. 1B). An additional expression vector, termed pILZ-Em, containing this zinc-inducible expression system with the same MCS and erythromycin resistance was also constructed from the plasmid pIL253 (Fig. 1B) (16). A BglII site was inserted into pIL253 by round PCR with the primers pIL-f and pIL-r (Table 1) in order to insert the BglII-SacI region from pCZ-Cm.

In order to assess the usefulness of this double inducible system, the structural gene of nisin, *nisA*, was cloned into plasmid pCZ-Cm under the control of *P_{czcD}* and transformed into NZ9800 (17). In this strain, the enzymes responsible for the maturation and modification of nisin are controlled by *P_{nisA}*. Comparison of the production of nisin in CDM medium (18, 19) with a constant concentration of nisin and various amounts of ZnSO₄ was performed (Fig. 4). We measured the production of nisin using an activity test against *L. lactis* NZ9000 (20). The activity assay clearly shows that nisin can be successfully expressed in the system in a tightly regulated fashion when the gene *nisA* is controlled by *P_{czcD}* and the modification enzymes are regulated by nisin (Fig. 4).

In our study, we introduced the streptococcal promoter *P_{czcD}* together with the gene coding for its regulatory protein, *SczA*, in *L. lactis*, yielding an effective zinc-regulated expression system, called Zirex. Our results clearly show that this system can effectively control the overexpression of proteins in response to mod-




FIG 4 Expression of nisin measured in terms of activity. In the control strain, NZ9800(pNZ-nisA) (20), *nisA* and the modification enzymes that process nisin are controlled by P_{nisA} . In NZ9800(pCZ-nisA), *nisA* was controlled by P_{czcD} . A constant concentration of nisin (5 ng/ml) was used to induce both strains. In NZ9800(pCZ-nisA), no activity was detected in the absence of Zn^{2+} , whereas increasing amounts of Zn^{2+} led to the production of nisin proportionally to the concentration used.

est and nontoxic zinc additions to the medium in *L. lactis*. The very low basal expression without inducer suggests that *SczA* is also expressed in *L. lactis* and tightly represses the system in the absence of zinc. Notably, *L. lactis* showed high tolerance to zinc in the millimolar range when induced in the exponential phase (6). Previously, a zinc-repressed expression system (P_{Zn} *zitR* promoter) was reported. It was based on the *L. lactis* *zit* operon, which encodes an emergency Zn^{2+} uptake ABC transporter (6). The presence of Zn^{2+} can repress the expression of the emergency Zn^{2+} uptake ABC transporter, which partly explains the high tolerance of *L. lactis* to Zn^{2+} . The initiation of the P_{Zn} *zitR* promoter is caused by the addition of a chelating agent, which reduces the available zinc in the medium, therefore activating the transcription of the emergency uptake system (6). A drawback of this system is that the induction based on the depletion of Zn^{2+} , which is achieved with the addition of EDTA, can hamper the overexpression of proteins or enzymes that require cations. The zinc-inducible system presented here constitutes, to our knowledge, the first zinc-inducible promoter developed for *L. lactis*. It can be extremely useful for the overproduction of enzymes such as lanthipeptide cyclases, which require Zn^{2+} to be active, or other metalloenzymes. This advantage makes the expression system presented in this paper a suitable candidate for the production of lanthipeptides (21). So far the nisin-inducible expression (NICE) system is the most widely used and potent protein expression system in *L. lactis*. Compared to the nisin-inducible system, the common drawbacks of other regulated expression systems found in *L. lactis* are their low expression level and/or high leakage (9). The zinc-inducible system presented here achieves a high expression level comparable to that of nisin (ca. 80%), which is higher than the expression level obtained with the P_{Zn} *zitR* promoter (20% of that achieved with nisin) (6). Moreover, we demonstrate that it is possible to combine both inducible promoters for the expression of different proteins at different times during cell growth. This can be a useful tool for the overexpression of proteins or the creation of controlled gene regulatory circuits in *L. lactis* and expands the toolbox available for this bacterium. Moreover, the plasmid described here could be directly applicable for use in other Gram-positive hosts, as is the case for the NICE system, although this has to be further investigated to assess the specific characteristics.

ACKNOWLEDGMENTS

We thank Tomas G. Kloosterman for the genomic DNA sample of *S. pneumoniae* D39 and helpful discussions.

D. Mu was supported by funding from the China Scholarship Council (no. 2010605032). M. Montalbán-López was supported by the NWO-ESF EuroSynbio program on a project called SynMod. Y. Masuda was funded by a grant from SNN (Province of Groningen) on project T2006, Lanthio-PEP-RUG.

REFERENCES

- de Ruyter PG, Kuipers OP, De Vos WM. 1996. Controlled gene expression systems for *Lactococcus lactis* with the food-grade inducer nisin. *Appl. Environ. Microbiol.* 62:3662–3667.
- Dieye Y, Usai S, Clier F, Gruss A, Piard J. 2001. Design of a protein-targeting system for lactic acid bacteria. *J. Bacteriol.* 183:4157–4166.
- van Asseldonk M, Simons A, Visser H, De Vos WM, Simons G. 1993. Cloning, nucleotide sequence, and regulatory analysis of the *Lactococcus lactis* *dnaJ* gene. *J. Bacteriol.* 175:1637–1644.
- Madsen SM, Arnau J, Vrang A, Givskov M, Israelsen H. 1999. Molecular characterization of the pH-inducible and growth phase-dependent promoter P170 of *Lactococcus lactis*. *Mol. Microbiol.* 32:75–87.
- Marugg JD, Meijer W, van Kranenburg R, Laverman P, Bruinenberg PG, De Vos WM. 1995. Medium-dependent regulation of proteinase gene expression in *Lactococcus lactis*: control of transcription initiation by specific dipeptides. *J. Bacteriol.* 177:2982–2989.
- Llull D, Poquet I. 2004. New expression system tightly controlled by zinc availability in *Lactococcus lactis*. *Appl. Environ. Microbiol.* 70:5398–5406.
- Kuipers OP, de Ruyter PG, Kleerebezem M, de Vos WM. 1998. Quorum sensing-controlled gene expression in lactic acid bacteria. *J. Biotechnol.* 64:15–21.
- Renye J, Jr, Somkuti G. 2010. Nisin-induced expression of pediocin in dairy lactic acid bacteria. *J. Appl. Microbiol.* 108:2142–2151.
- Kuipers OP, de Ruyter PG, Kleerebezem M, de Vos WM. 1997. Controlled overproduction of proteins by lactic acid bacteria. *Trends Biotechnol.* 15:135–140.
- Kloosterman TG, Der Kooi-Pol V, Magdalena M, Bijlsma JJ, Kuipers OP. 2007. The novel transcriptional regulator *SczA* mediates protection against Zn^{2+} stress by activation of the Zn^{2+} -resistance gene *czcD* in *Streptococcus pneumoniae*. *Mol. Microbiol.* 65:1049–1063.
- Eberhardt A, Wu LJ, Errington J, Vollmer W, Veening J. 2009. Cellular localization of choline-utilization proteins in *Streptococcus pneumoniae* using novel fluorescent reporter systems. *Mol. Microbiol.* 74:395–408.
- Lanie JA, Ng WL, Kazmierczak KM, Andrzejewski TM, Davidsen TM, Wayne KJ, Tettelin H, Glass JI, Winkler ME. 2007. Genome sequence of Avery's virulent serotype 2 strain D39 of *Streptococcus pneumoniae* and comparison with that of unencapsulated laboratory strain R6. *J. Bacteriol.* 189:38–51.
- Higuchi R, Krummel B, Saiki R. 1988. A general method of *in vitro* preparation and specific mutagenesis of DNA fragments: study of protein and DNA interactions. *Nucleic Acids Res.* 16:7351–7367.
- Holo H, Nes I. 1995. Transformation of *Lactococcus* by electroporation. *Methods Mol. Biol.* 47:195–199.
- Solopova A, Bachmann H, Teusink B, Kok J, Neves AR, Kuipers OP. 2012. A specific mutation in the promoter region of the silent *cel* cluster accounts for the appearance of lactose-utilizing *Lactococcus lactis* MG1363. *Appl. Environ. Microbiol.* 78:5612–5621.

16. Simon D, Chopin A. 1988. Construction of a vector plasmid family and its use for molecular cloning in *Streptococcus lactis*. *Biochimie* 70:559–566.
17. Kuipers OP, Beerthuyzen MM, Siezen RJ, De Vos WM. 1993. Characterization of the nisin gene cluster *nisABTCIPR* of *Lactococcus lactis*: requirement of expression of the *nisA* and *nisI* genes for development of immunity. *Eur. J. Biochem.* 216:281–291.
18. Otto R, Brink B, Veldkamp H, Konings WN. 1983. The relation between growth rate and electrochemical proton gradient of *Streptococcus cremoris*. *FEMS Microbiol. Lett.* 16:69–74.
19. Poolman B, Konings WN. 1988. Relation of growth of *Streptococcus lactis* and *Streptococcus cremoris* to amino acid transport. *J. Bacteriol.* 170:700–707.
20. van Heel AJ, Mu D, Montalbán-López M, Hendriks D, Kuipers OP. 1 February 2013. Designing and producing modified, new-to-nature, peptides with antimicrobial activity by use of a combination of various lantibiotic modification enzymes. *ACS Synth. Biol.* doi:10.1021/sb3001084.
21. Li B, Yu JPJ, Brunzelle JS, Moll GN, Van der Donk WA, Nair SK. 2006. Structure and mechanism of the lantibiotic cyclase involved in nisin biosynthesis. *Science* 311:1464–1467.